



## Risk of Falling

You may be at **risk for falling**. Both Primary Periodic Paralysis (PPP) and treatment with KEVEYIS for PPP may increase your risk.<sup>1</sup> This brochure can help you **talk with your healthcare team** about how to best reduce the risk of falling.

**Please see Important Safety Information on back  
and Full Prescribing Information in pocket.**



**KEVEYIS<sup>®</sup>**  
dichlorphenamide 50 mg tablets

# Feeling more likely to fall?

The risk of falling can be due to many different factors, like the environment around you, a medical condition, or certain medications like KEVEYIS® (dichlorphenamide).<sup>1-3</sup>

Here are some strategies that may help reduce your risk of falling.

---

This information is provided by **Strongbridge Biopharma** as part of our commitment to providing patients a wide range of services and support. **For more information, call 844-538-3947.**

---

*The content included herein is not intended to be a substitute for professional medical advice or treatment. Always seek the advice of your doctor or other qualified healthcare provider with any questions you may have about a medical condition.*



## Search for the source.

It could be due to the medication you're taking, so talk to your doctor about your KEVEYIS dose.<sup>1</sup>

---



## Wear comfortable shoes.

Think about what you have on your feet and consider where you're going to be walking. Wear properly fitting shoes with nonskid soles and avoid hazards at home.<sup>2,4</sup>

---



## Stand strong.

Ask your doctor about the potential benefits of implementing a strength and balance exercise regimen. The conversation should occur before you begin to exercise.<sup>4</sup>

---



## Don't be afraid to ask for help.

There's no shame in asking for a hand to hold or a shoulder to lean on. Those around you will understand.

---

**When using these tips, keep in mind that everyone is different. Talk to your doctor and find out what's best for you.**

---



**KEVEYIS**<sup>®</sup>  
dichlorphenamide 50 mg tablets

## What is KEVEYIS?

KEVEYIS (pronounced keh-VAY-iss) (dichlorphenamide) is a prescription drug used to treat primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and other similar diseases.

## Important Safety Information

### What should you tell your healthcare provider before taking KEVEYIS?

Tell your doctor if you are allergic to dichlorphenamide or other sulfa drugs; if you take high doses of aspirin, or if you have lung or liver disease; if you are pregnant, plan to become pregnant, or are breastfeeding or plan to breastfeed.

### What should you know about KEVEYIS?

- Severe allergic and other reactions have happened with sulfonamides (drugs such as KEVEYIS) and have sometimes been fatal. Stop taking KEVEYIS at the first sign of skin rash, swelling, difficulty breathing, or any other unexpected side effect or reaction, and call 911 right away.
- Tell your healthcare provider if you take aspirin or if another healthcare provider instructs you to begin taking aspirin. High doses of aspirin should not be taken with KEVEYIS.
- KEVEYIS can cause your body to lose potassium, which can lead to heart problems. Your healthcare provider will measure the potassium levels in your blood before you start treatment and at certain times during treatment.
- Tell your healthcare provider about all other prescription and over-the-counter medicines you take, including supplements, as some medicines can interact with KEVEYIS.
- While taking KEVEYIS, your body may produce too much acid or may not be able to remove acid from the body. Your healthcare provider may run tests on a regular basis to check for signs of acid buildup.
- KEVEYIS may increase your risk of falling. Tell your healthcare provider right away if you experience a fall while taking KEVEYIS.
- The risks of falls and acid buildup are greater in elderly patients.
- It is not known whether KEVEYIS is safe or effective for people younger than 18 years of age.

### What are the most common side effects with KEVEYIS?

The most common side effects are a feeling of numbness, tingling or burning ("pins and needles") in the toes, feet, hands or fingers; trouble with memory or thinking; feeling confused; and unpleasant taste in the mouth.

These are not all of the possible side effects of KEVEYIS. Call your healthcare provider for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

### Please see Full Prescribing Information in the pocket.

**References.** **1.** KEVEYIS [package insert]. Feasterville-Trevose, PA: Strongbridge Biopharma; 2017. **2.** Fall prevention: simple tips to prevent falls. Mayo Clinic. <https://www.mayoclinic.org/healthy-lifestyle/healthy-aging/in-depth/fall-prevention/art-20047358>. Accessed July 20, 2018. **3.** Statland JM, Fontaine B, Hanna MG, et al. Review of the diagnosis and treatment of periodic paralysis. *Muscle Nerve*. 2018;57:522-530. **4.** Talking about fall prevention with your patients. Centers for Disease Control and Prevention. <https://www.cdc.gov/steady/pdf/STEADI-FactSheet-TalkingWPpatients-508.pdf>. Accessed July 20, 2018.

900 Northbrook Drive, Suite 200  
Trevose, PA 19053 United States  
[www.keveyis.com](http://www.keveyis.com)

© 2018 Strongbridge Biopharma plc

STRONGBRIDGE BIOPHARMA® is a registered trademark of Strongbridge Biopharma plc. companies, which include Strongbridge Ireland Limited and Strongbridge U.S. Inc.

KEVEYIS® is a registered trademark licensed exclusively in the US to Strongbridge Biopharma plc.

KEV-0375 07/2018

**KEVEYIS**<sup>®</sup>  
dichlorphenamide 50 mg tablets

**STRONGBRIDGE**  
BIOPHARMA

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use KEVEYIS® safely and effectively. See full prescribing information for KEVEYIS®.

**KEVEYIS® (dichlorphenamide) tablets, for oral use**

**Initial U.S. Approval: 1958**

### RECENT MAJOR CHANGES

Indications and Usage: treatment of primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and related variants (1) 8/2015  
Dosage and Administration (2) 8/2015  
Warnings and Precautions (5.1, 5.4, 5.5) 8/2015

### INDICATIONS AND USAGE

KEVEYIS is an oral carbonic anhydrase inhibitor indicated for the treatment of primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and related variants (1)

### DOSAGE AND ADMINISTRATION

- Initial dose: 50 mg twice daily (2)
- Titrate dose based on individual response (2)
- The maximum recommended dose is 200 mg daily (2)

### DOSAGE FORMS AND STRENGTHS

Tablets: 50 mg (3)

### CONTRAINDICATIONS

- Hepatic insufficiency (4)
- Severe pulmonary obstruction (4)
- Hypersensitivity to dichlorphenamide or other sulfonamides (4)
- Concomitant use with high dose aspirin (4)

### WARNINGS AND PRECAUTIONS

- Hypersensitivity / Anaphylaxis / Idiosyncratic reactions: discontinue KEVEYIS at the first appearance of skin rash or any sign of immune-mediated or idiosyncratic adverse reaction (5.1)
- Hypokalemia: baseline and periodic measurement of serum potassium are recommended; if hypokalemia develops or persists, consider reducing the dose or discontinuing KEVEYIS (5.3)
- Metabolic acidosis: baseline and periodic measurement of serum bicarbonate are recommended; if metabolic acidosis develops or persists, consider reducing the dose or discontinuing KEVEYIS(5.4)
- Falls: consider reducing the dose or discontinuing KEVEYIS in patients who experience falls (5.5)

### ADVERSE REACTIONS

Most common adverse reactions (incidence at least 10% and greater than placebo) include paresthesias, cognitive disorder, dysgeusia, and confusional state (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Strongbridge Biopharma at 1-855-324-8912, or the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

Aspirin: Anorexia, tachypnea, lethargy, and coma have been reported with concomitant use of dichlorphenamide and high-dose aspirin. The concomitant use of KEVEYIS and high dose aspirin is contraindicated. KEVEYIS should be used with caution in patients receiving low dose aspirin (4, 5.2, 7.1).

### USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm. (8.1)

**See 17 for PATIENT COUNSELING INFORMATION**

**Revised: 01/2017**

## FULL PRESCRIBING INFORMATION: CONTENTS\*

- 1 INDICATIONS AND USAGE**
- 2 DOSAGE AND ADMINISTRATION**
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
  - 5.1 Hypersensitivity / Anaphylaxis / Idiosyncratic Reactions
  - 5.2 Concomitant Use of Aspirin
  - 5.3 Hypokalemia
  - 5.4 Metabolic Acidosis
  - 5.5 Falls
- 6 ADVERSE REACTIONS**
  - 6.1 Clinical Trials Experience
  - 6.2 Postmarketing Experience
- 7 DRUG INTERACTIONS**
  - 7.1 Aspirin and Salicylates

- 8 USE IN SPECIFIC POPULATIONS**
  - 8.1 Pregnancy
  - 8.3 Nursing Mothers
  - 8.4 Pediatric Use
  - 8.5 Geriatric Use
- 10 OVERDOSAGE**
- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
  - 12.1 Mechanism of Action
  - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
  - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

KEVEYIS is indicated for the treatment of primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and related variants.

### 2 DOSAGE AND ADMINISTRATION

Initiate dosing at 50 mg twice daily. The initial dose may be increased or decreased based on individual response, at weekly intervals (or sooner in case of adverse reaction). The maximum recommended total daily dose is 200 mg.

Primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and related variants are a heterogeneous group of conditions, for which the response to KEVEYIS may vary. Therefore, prescribers should evaluate the patient's response to KEVEYIS after 2 months of treatment to decide whether KEVEYIS should be continued.

### 3 DOSAGE FORMS AND STRENGTHS

Round, white tablets, scored on one side, engraved with “D” above the score and “50” below the score, the other side is plain, 50 mg each.

### 4 CONTRAINDICATIONS

KEVEYIS is contraindicated in the following circumstances:

- Hypersensitivity to dichlorphenamide or other sulfonamides [*see Warnings and Precautions (5.1)*]
- Concomitant use of KEVEYIS and high dose aspirin [*see Warnings and Precautions (5.2)*]
- Severe pulmonary disease, limiting compensation to metabolic acidosis caused by KEVEYIS [*see Warnings and Precautions (5.4)*]
- Hepatic insufficiency: KEVEYIS may aggravate hepatic encephalopathy.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hypersensitivity / Anaphylaxis / Idiosyncratic Reactions

Fatalities associated with the administration of sulfonamides have occurred due to adverse reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis, fulminant hepatic necrosis, agranulocytosis, aplastic anemia and other blood dyscrasias. Pulmonary involvement can occur in isolation or as part of a systemic reaction.

KEVEYIS should be discontinued at the first appearance of skin rash or any sign of immune-mediated or idiosyncratic adverse reaction.

## **5.2 Concomitant Use of Aspirin**

Anorexia, tachypnea, lethargy, and coma have been reported with concomitant use of dichlorophenamide and high-dose aspirin. The concomitant use of KEVEYIS and high dose aspirin is contraindicated. KEVEYIS should be used with caution in patients receiving low dose aspirin.

## **5.3 Hypokalemia**

KEVEYIS increases potassium excretion and can cause hypokalemia. The risk of hypokalemia is greater when KEVEYIS is used in patients with conditions associated with hypokalemia (e.g., adrenocortical insufficiency, hyperchloremic metabolic acidosis, or respiratory acidosis), and in patients receiving other drugs that may cause hypokalemia (e.g., loop diuretics, thiazide diuretics, laxatives, antifungals, penicillin, and theophylline).

Baseline and periodic measurement of serum potassium during KEVEYIS treatment are recommended.

If hypokalemia develops or persists, consideration should be given to reducing the dose or discontinuing KEVEYIS.

## **5.4 Metabolic Acidosis**

KEVEYIS can cause hyperchloremic non-anion gap metabolic acidosis. Concomitant use of KEVEYIS with other drugs that cause metabolic acidosis may increase the severity of metabolic acidosis.

Baseline and periodic measurement of serum bicarbonate during KEVEYIS treatment are recommended.

If metabolic acidosis develops or persists, consideration should be given to reducing the dose or discontinuing KEVEYIS.

## **5.5 Falls**

KEVEYIS increases the risk of falls. The risk of falls is greater in the elderly and with higher doses of KEVEYIS. Consider dose reduction or discontinuation of KEVEYIS in patients who experience falls while treated with KEVEYIS.

## 6 ADVERSE REACTIONS

The following serious adverse reactions are described elsewhere in labeling:

- Hypersensitivity / Anaphylaxis / Idiosyncratic reactions [see *Warnings and Precautions (5.1)*]
- Hypokalemia [see *Warnings and Precautions (5.3)*]
- Metabolic Acidosis [see *Warnings and Precautions (5.4)*]
- Falls [see *Warnings and Precautions (5.5)*]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In a 9-week randomized controlled trial in adults with hyperkalemic or hypokalemic periodic paralysis (Study 1), the most common adverse reactions in patients treated with KEVEYIS, with rates greater than placebo, were paresthesia, cognitive disorder, dysgeusia, and confusional state. The mean dose of KEVEYIS was 94 mg/day in patients with hypokalemic periodic paralysis and 82 mg/day in patients with hyperkalemic periodic paralysis.

Table 1 lists the incidence of adverse reactions that occurred in  $\geq 5\%$  of patients treated with KEVEYIS and more commonly than in patients treated with placebo in Study 1.

**Table 1: Adverse Reactions in Patients Treated with KEVEYIS with Incidence  $\geq 5\%$  and more common than in Patients Treated with Placebo in Study 1**

	<b>Adverse Reaction</b>	<b>KEVEYIS</b> N = 36 (%)	<b>Placebo</b> N = 29 (%)
Nervous system disorders	Paresthesia	44	14
	Cognitive disorder <sup>1</sup>	14	7
	Dysgeusia	14	0
	Confusional state	11	0
	Headache	8	7
	Hypoesthesia	8	0
	Lethargy	8	0
	Dizziness	6	0
Gastrointestinal disorders	Diarrhea	6	3
	Nausea	6	0
General disorders and administration site conditions	Fatigue	8	0
	Malaise	6	0
Investigations	Weight decreased	6	0
Musculoskeletal and connective tissue disorders	Muscle spasms	8	0
	Arthralgia	6	3
	Muscle twitching	6	0
Respiratory	Dyspnea	6	0
	Pharyngolaryngeal pain	6	0
Skin	Rash	8	0
	Pruritus	6	0



<sup>1</sup> Cognitive disorder combined cases with the preferred terms of *cognitive disorder, disturbance in attention, and mental impairment*.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during postapproval use of dichlorphenamide. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following are adverse reactions which have been reported for dichlorphenamide that were serious adverse events or are not reported in the previous section of labeling [*see Clinical Trials Experience (6.1)*]: amnesia, cardiac failure, condition aggravated, convulsion, fetal death, hallucination, nephrolithiasis, pancytopenia, psychotic disorder, renal tubular necrosis, stupor, syncope, tremor.

## **7 DRUG INTERACTIONS**

### **7.1 Aspirin and Salicylates**

KEVEYIS may cause an elevation in salicylate levels in patients receiving aspirin. Anorexia, tachypnea, lethargy, and coma have been reported with concomitant use of dichlorphenamide and high-dose aspirin.

Concomitant use of KEVEYIS and high dose aspirin is contraindicated. KEVEYIS should be used with caution in patients receiving low dose aspirin. [*see Contraindications (4) and Warnings and Precautions (5.2)*]

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

Pregnancy Category C.

There are no adequate and well-controlled studies in pregnant women. Teratogenic effects (fetal limb reduction defects) were reported following oral administration of dichlorphenamide to pregnant rats during organogenesis at 350 mg/kg, or 17 times the maximum recommended human dose (200 mg/day) on a body surface area (mg/m<sup>2</sup>) basis. A no-effect dose has not been established. KEVEYIS should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

### 8.3 Nursing Mothers

It is not known whether dichlorphenamide is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when dichlorphenamide is administered to a nursing woman.

### 8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

### 8.5 Geriatric Use

The risk of falls and of metabolic acidosis are greater in elderly patients.

## 10 OVERDOSAGE


Symptoms of overdosage or toxicity may include drowsiness, anorexia, nausea, vomiting, dizziness, paresthesias, ataxia, tremor, and tinnitus.

In the event of overdosage, induce emesis or perform gastric lavage. The electrolyte disturbance most likely to be encountered from overdosage is hyperchloremic acidosis.

## 11 DESCRIPTION

KEVEYIS (dichlorphenamide) tablets is an oral carbonic anhydrase inhibitor. Dichlorphenamide, a dichlorinated benzenedisulfonamide, is known chemically as 4, 5-dichloro-1,3-benzenedisulfonamide.

Its empirical formula is  $C_6H_4Cl_2N_2O_4S_2$  and its structural formula is:



Dichlorphenamide USP is a white or practically white, crystalline compound with a molecular weight of 305.16. It is very slightly soluble in water but soluble in dilute solutions of sodium carbonate and sodium hydroxide. Dilute alkaline solutions of dichlorphenamide are stable at room temperature.

KEVEYIS (dichlorphenamide) tablets is supplied as tablets, for oral administration, each containing 50 mg dichlorphenamide. Inactive ingredients are lactose monohydrate, magnesium stearate and pregelatinized maize starch.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Dichlorphenamide is a carbonic anhydrase inhibitor. However, the precise mechanism by which dichlorphenamide exerts its therapeutic effects in patients with periodic paralysis is unknown.

### **12.3 Pharmacokinetics**

The pharmacokinetic properties of dichlorphenamide after oral absorption are not known.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### **Carcinogenesis**

Studies to assess the carcinogenic potential of dichlorphenamide have not been conducted.

#### **Mutagenesis**

Studies to assess the genotoxicity of dichlorphenamide have not been conducted.

#### **Impairment of Fertility**

Studies to assess the effects of dichlorphenamide on fertility have not been conducted.

## **14 CLINICAL STUDIES**

The efficacy of KEVEYIS was evaluated in two clinical studies, Study 1 and Study 2.

### Study 1

Study 1 was a 9-week, double blind, placebo-controlled multi-center study. Study 1 consisted of two substudies: a substudy in patients with hypokalemic periodic paralysis (n=44), and a substudy in patients with hyperkalemic periodic paralysis (n=21). The primary efficacy endpoint in both substudies was the average number of self-reported attacks of muscle weakness per week over the final 8 weeks of the trial. Withdrawal from the study for acute severe worsening was also assessed as an endpoint.

In Study 1, the dose of KEVEYIS was 50 mg b.i.d. for treatment-naïve patients. Patients already on dichlorphenamide prior to the study continued on the same dose while on KEVEYIS during

the study. In patients taking acetazolamide prior to the study, the dose of KEVEYIS was set at 20% of the acetazolamide dose. Dose reduction for tolerability was permitted.

#### *Hypokalemic Periodic Paralysis Substudy of Study 1*

In the hypokalemic periodic paralysis substudy, median age of patients was 45 years and 73% of patients were male. Patients treated with KEVEYIS (n=24) had 2.2 fewer attacks per week than patients (n=20) treated with placebo (p=0.02). None of the patients randomized to KEVEYIS reached the endpoint of acute worsening, vs. five patients randomized to placebo. The mean dose of KEVEYIS at Week 9 was 94 mg/day.

#### *Hyperkalemic Periodic Paralysis Substudy of Study 1*

In the Hyperkalemic Periodic Paralysis substudy, median age of patients was 43 years and 43% of patients were male. During the double-blind treatment period, patients treated with KEVEYIS (n=12) had 3.9 fewer attacks per week than patients (n=9) treated with placebo (p=0.08). None of the patients randomized to KEVEYIS reached the endpoint of acute worsening, vs. two patients randomized to placebo. The mean dose of KEVEYIS at Week 9 was 82 mg/day.

#### Study 2

Study 2 was a 35-week, double blind, placebo-controlled, multi-center, two-period crossover study. Study 2 also consisted of two substudies: a substudy in a substudy in patients with hypokalemic periodic paralysis (n=42), and a substudy in patients with hyperkalemic periodic paralysis (n=31), including patients with Paramyotonia Congenita. The primary endpoint in the hypokalemic periodic paralysis substudy was the incidence of acute intolerable worsening (based on attack frequency or severity) necessitating withdrawal. The primary endpoint in the hyperkalemic periodic paralysis substudy was the average number of self-reported attacks of muscle weakness per week. Dosing was determined similarly to Study 1.

#### *Hypokalemic Periodic Paralysis Substudy of Study 2*

In the hypokalemic periodic paralysis substudy, mean age of patients was 38 years and 79% of patients were male. Acute intolerable worsening was observed in 2 patients on KEVEYIS vs. 11 patients on placebo (p=0.02). The mean dose of KEVEYIS at the end of the study was 96 mg/day.

#### *Hyperkalemic Periodic Paralysis Substudy of Study 2*

In the hyperkalemic periodic paralysis substudy, mean age of patients was 37 years and 79% of patients were male. Patients treated had 2.3 fewer attacks per week on KEVEYIS than on placebo (p=0.006). The mean dose of KEVEYIS at the end of the study was 73 mg/day.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Each KEVEYIS (dichlorphenamide) tablet, 50 mg is round, white, scored on one side, engraved with “D” above the score and “50” below the score. The other side is plain.

KEVEYIS (dichlorphenamide) tablets are supplied as follows:

Bottles of 100 .....NDC 71090-001-01

Store at 20° to 25° C (68° to 77° F) [See USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

### Worsening of Symptoms

Advise patients to notify their physician if they experience worsening of symptoms of periodic paralysis.

### Driving and Operating Machinery

KEVEYIS may cause drowsiness/fatigue in some patients. Caution patients on the potential for impaired ability to drive and operate machinery.

Mfd. by: Taro Pharmaceutical Industries Ltd., Haifa Bay, Israel 2624761

Dist. by: Strongbridge US Inc., Trevose, PA 19053

KEVEYIS<sup>®</sup> is a licensed trademark of Strongbridge Biopharma plc.

STRONGBRIDGE BIOPHARMA<sup>™</sup> is a trademark of Strongbridge Biopharma plc.

Strongbridge Biopharma plc is the parent company of Strongbridge US Inc.

Revised: January 2017

21061-0117-0

